

# An introduction to shape and topology optimization

Éric Bonnetier\* and Charles Dapogny†

\* Institut Fourier, Université Grenoble-Alpes, Grenoble, France

† CNRS & Laboratoire Jean Kuntzmann, Université Grenoble-Alpes, Grenoble, France

Fall, 2020


## Foreword: geometric shape optimization

We have seen how to optimize shapes when they are **parametrized**:


$$\min_h J(h) \text{ s.t. } C(h) \leq 0,$$

where the **design variable**  $h$  may be:

- A set of parameters in a finite-dimensional space (thickness, etc.);
- A function  $h$  in a suitable, infinite dimensional vector (Banach) space.



Description of a mechanical part via the control points of a CAD model.



Parametrization of a plate with cross-section  $S$  via the thickness function  $h : S \rightarrow \mathbb{R}$ .

## Foreword: geometric shape optimization (II)

### Assets:

- In the considered examples, the state  $u_h$  lives in a fixed computational domain, which greatly simplifies the calculation of **derivatives with respect to the design**.
- Efficient methods from mathematical programming (optimization routines, etc.) are readily available in this context.

### Drawbacks:

- This induces a strong **bias** in the sought shapes.
- It may be very difficult, and in practice cumbersome, to find which are the relevant parameters  $h$  of shapes.

⇒ It is often desirable to formulate shape optimization problems in terms of the **geometry** of shapes  $\Omega$ :

$$\min J(\Omega) \text{ s.t. } C(\Omega) \leq 0.$$

## Part III

# Geometric optimization problems


- 1 The method of Hadamard and shape derivatives
- 2 Shape derivatives of PDE-constrained functionals: the rigorous way, using Eulerian and material derivatives
- 3 C ea's method for calculating shape derivatives
- 4 Numerical aspects of geometric methods
- 5 The level set method for shape optimization

## Preliminary notations

Let  $\Omega$  be a bounded domain in  $\mathbb{R}^d$ ;

- $\partial\Omega$  is the **boundary** of  $\Omega$ ;
- $n : \partial\Omega \rightarrow \mathbb{R}^d$  denotes the **unit normal vector** to  $\partial\Omega$ , pointing outward  $\Omega$ ;
- The domain  $\Omega$  is called **Lipschitz** (resp. of class  $\mathcal{C}^k$ ) if

*“Near every point  $x \in \partial\Omega$ ,  $\Omega$  resembles the lower part of the graph of a Lipschitz function (resp. of a  $\mathcal{C}^k$  function).”*




In a neighborhood  $U$  of each point  $x \in \partial\Omega$ ,  $\Omega$  “looks like” the lower part of the graph of some (Lipschitz or  $\mathcal{C}^k$ ) function  $\hat{x} \mapsto f(\hat{x})$  defined for suitable  $(d - 1)$ -dimensional coordinates.

**Hadamard's boundary variation method** describes variations of a reference, bounded Lipschitz domain  $\Omega$  of the form:

$$\Omega \mapsto \Omega_\theta := (\text{Id} + \theta)(\Omega),$$

for "small" vector fields  $\theta \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$ .



## Lemma 1.

For  $\theta \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$  with norm  $\|\theta\|_{W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)} < 1$ , the mapping  $(\text{Id} + \theta)$  is a Lipschitz diffeomorphism.

## Definition 1.

Given a bounded Lipschitz domain  $\Omega$ , a function  $\Omega \mapsto J(\Omega) \in \mathbb{R}$  is **shape differentiable** at  $\Omega$  if the mapping

$$W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d) \ni \theta \mapsto J(\Omega_\theta)$$

is Fréchet-differentiable at 0, i.e. the following expansion holds in the vicinity of 0:

$$J(\Omega_\theta) = J(\Omega) + J'(\Omega)(\theta) + o(\theta), \text{ where } \frac{o(\theta)}{\|\theta\|_{W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)}} \xrightarrow{\theta \rightarrow 0} 0..$$

The linear mapping  $\theta \mapsto J'(\Omega)(\theta)$  is the **shape derivative** of  $J$  at  $\Omega$ .

**Remark** Other spaces are often used in place of  $W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$ , made of **more regular** deformation fields  $\theta$ , e.g.:

$$C^{k,\infty}(\mathbb{R}^d, \mathbb{R}^d) := \left\{ \theta : \mathbb{R}^d \rightarrow \mathbb{R}^d \text{ of class } C^k, \sup_{|\alpha| \leq k} \sup_{x \in \mathbb{R}^d} |\partial^\alpha \theta(x)| < \infty \right\}.$$

## Theorem 2.

Let  $\Omega \subset \mathbb{R}^d$  be a bounded *Lipschitz* domain, and let  $f \in W^{1,1}(\mathbb{R}^d)$  be a *fixed* function. Consider the functional:


$$J(\Omega) = \int_{\Omega} f(x) \, dx;$$

then  $J(\Omega)$  is shape differentiable at  $\Omega$  and its shape derivative is:

$$\forall \theta \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d), \quad J'(\Omega)(\theta) = \int_{\partial\Omega} f(\theta \cdot n) \, ds.$$



## First examples of shape derivatives (II)



*Intuition:*  $f$  takes negative (resp. positive) values on the blue (resp. red) part of the boundary  $\partial\Omega$ . The value  $J(\Omega_\theta)$  is minimized from  $J(\Omega)$  by adding the blue area, (i.e.  $\theta \cdot n > 0$  where  $f < 0$ ), and by removing the red area ( $\theta \cdot n < 0$  where  $f > 0$ ), weighted by  $f$ .

## First examples of shape derivatives (III)

### Remarks:

- This result is a particular case of the **Transport** (or **Reynolds**) **theorem**, used to derive the equations of motion from conservation principles in fluid mechanics (see the Appendix in **Lecture 1**).
- It allows to calculate the shape derivative of the **volume** functional

$$\text{Vol}(\Omega) = \int_{\Omega} 1 \, dx;$$

Indeed, it holds:

$$\forall \theta \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d), \quad \text{Vol}'(\Omega)(\theta) = \int_{\partial\Omega} \theta \cdot n \, ds = \int_{\Omega} \text{div} \theta \, dx.$$

In particular, if  $\text{div} \theta = 0$ , the volume is unchanged (at first order) when  $\Omega$  is perturbed by  $\theta$ .

## First examples of shape derivatives (IV)

Proof: The formula proceeds from a **change of variables** in volume integrals:

$$J(\Omega_\theta) = \int_{(\text{Id}+\theta)(\Omega)} f(x) \, dx = \int_{\Omega} |\det(\text{Id} + \nabla\theta)| f \circ (\text{Id} + \theta) \, dx.$$

- The mapping  $\theta \mapsto \det(\text{Id} + \nabla\theta)$  is Fréchet differentiable, and:

$$\det(\text{Id} + \nabla\theta) = 1 + \text{div}\theta + o(\theta), \text{ where } \frac{o(\theta)}{\|\theta\|_{W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)}} \xrightarrow{\theta \rightarrow 0} 0.$$

- If  $f \in W^{1,1}(\mathbb{R}^d)$ ,  $\theta \mapsto f \circ (\text{Id} + \theta)$  is also Fréchet differentiable and:

$$f \circ (\text{Id} + \theta) = f + \nabla f \cdot \theta + o(\theta).$$

- Combining those three identities and **Green's formula** leads to the result. □

**Remark:** This idea of

- ① Using the change of variables  $\Omega \rightarrow (\text{Id} + \theta)(\Omega)$  to transport all integrals on the reference domain  $\Omega$ ,
- ② Differentiating with respect to the deformation  $\theta$ ,

is the “standard” way to calculate shape derivatives.

### Theorem 3.

Let  $\Omega \subset \mathbb{R}^d$  be a bounded domain of class  $\mathcal{C}^2$ , and let  $g \in W^{2,1}(\mathbb{R}^d)$  be a **fixed** function. Consider the functional:

$$J(\Omega) = \int_{\partial\Omega} g(x) \, ds;$$

then  $J(\Omega)$  is shape differentiable at  $\Omega$  when deformations  $\theta$  are chosen in

$$\mathcal{C}^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d) := \mathcal{C}^1(\mathbb{R}^d, \mathbb{R}^d) \cap W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d),$$

and the shape derivative is:


$$J'(\Omega)(\theta) = \int_{\partial\Omega} \left( \frac{\partial g}{\partial n} + \kappa g \right) (\theta \cdot n) \, ds,$$

where  $\kappa$  is the **mean curvature** of  $\partial\Omega$ .

**Example:** The shape derivative of the **perimeter**  $\text{Per}(\Omega) = \int_{\partial\Omega} 1 \, ds$  is:

$$\text{Per}'(\Omega)(\theta) = \int_{\partial\Omega} \kappa (\theta \cdot n) \, ds.$$


## First examples of shape derivatives (VI)



Intuition:  $\theta = -\kappa n$  is a *descent direction* for  $\text{Per}(\Omega)$ : it is reduced by smearing the *bumps* of  $\partial\Omega$  (i.e.  $\theta \cdot n < 0$  when  $\kappa > 0$ ), and sealing its *holes* (i.e.  $\theta \cdot n > 0$  when  $\kappa < 0$ ).

## Structure of shape derivatives (I)

**Idea:** The shape derivative  $J'(\Omega)(\theta)$  of a “regular” functional  $\Omega \mapsto J(\Omega)$  only depends on the normal component  $\theta \cdot n$  of the vector field  $\theta$ .



At first order, a **tangential** vector field  $\theta$ , (i.e.  $\theta \cdot n = 0$ ) only results in a **convection** of the shape  $\Omega$ , and it is expected that  $J'(\Omega)(\theta) = 0$ .

## Lemma 4.

Let  $\Omega$  be a domain of class  $C^1$ . Assume that the mapping

$$C^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d) \ni \theta \mapsto J(\Omega_\theta) \in \mathbb{R}$$

is of class  $C^1$ . Then, for any vector field  $\theta \in C^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$  such that  $\theta \cdot n = 0$  on  $\partial\Omega$ , one has:  $J'(\Omega)(\theta) = 0$ .

## Corollary 5.

Under the same hypotheses, if  $\theta_1, \theta_2 \in C^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$  have the same normal component, i.e.  $\theta_1 \cdot n = \theta_2 \cdot n$  on  $\partial\Omega$ , then:

$$J'(\Omega)(\theta_1) = J'(\Omega)(\theta_2).$$

## Structure of shape derivatives (III)

- Actually, the shape derivatives of “many” integral objective functionals  $J(\Omega)$  can be put under the **surface form**:

$$J'(\Omega)(\theta) = \int_{\partial\Omega} v_{\Omega} (\theta \cdot n) \, ds,$$

where the scalar field  $v_{\Omega} : \partial\Omega \rightarrow \mathbb{R}$  depends on  $J$  and on the current shape  $\Omega$ .

- This structure lends itself to the calculation of a **descent direction**: letting  $\theta = -tv_{\Omega}n$ , for a small enough **descent step**  $t > 0$  in the definition of shape derivatives yields:

$$J(\Omega_{t\theta}) = J(\Omega) - t \int_{\partial\Omega} v_{\Omega}^2 \, ds + o(t) < J(\Omega).$$

- We shall return to this issue during our study of numerical algorithms.



## Part III

# Geometric optimization problems

- 1 The method of Hadamard and shape derivatives
- 2 Shape derivatives of PDE-constrained functionals: the rigorous way, using Eulerian and material derivatives**
- 3 C ea's method for calculating shape derivatives
- 4 Numerical aspects of geometric methods
- 5 The level set method for shape optimization

## Shape derivatives of PDE constrained functionals

- Hitherto, we have studied the shape derivatives of functionals of the form

$$F_1(\Omega) = \int_{\Omega} f(x) \, dx, \text{ and } F_2(\Omega) = \int_{\partial\Omega} g(x) \, ds,$$

where  $f, g : \mathbb{R}^d \rightarrow \mathbb{R}$  are given, smooth enough functions.

- We now intend to consider functions of the form

$$J_1(\Omega) = \int_{\Omega} j(u_{\Omega}(x)) \, dx, \text{ or } J_2(\Omega) = \int_{\partial\Omega} k(u_{\Omega}(x)) \, ds,$$

where  $j, k : \mathbb{R} \rightarrow \mathbb{R}$  are given, smooth enough functions, and  $u_{\Omega} : \Omega \rightarrow \mathbb{R}$  is **the solution to a PDE posed on  $\Omega$** .

- Doing so elaborates on the techniques from optimal control theory that we have seen in the parametric optimization context.

## The considered framework

- For simplicity, we rely on the simplified model of the Laplace equation with Dirichlet boundary conditions: the state  $u_\Omega$  is solution to:

$$\begin{cases} -\Delta u_\Omega = f & \text{in } \Omega \\ u_\Omega = 0 & \text{on } \partial\Omega, \end{cases}$$

for a smooth enough source  $f : \mathbb{R}^d \rightarrow \mathbb{R}$ .

- The associated variational formulation reads:

$$\forall v \in H_0^1(\Omega), \quad \int_{\Omega} \nabla u_\Omega \cdot \nabla v \, dx = \int_{\Omega} f v \, dx.$$

- In this setting:
  - ① We calculate the “derivative” of the state  $\Omega \mapsto u_\Omega$  in a sense to be defined.
  - ② We infer the shape derivative of a shape functional of the form:

$$J(\Omega) = \int_{\Omega} j(u_\Omega) \, dx,$$

where  $j : \mathbb{R} \rightarrow \mathbb{R}$  is a “smooth enough” function.


## Eulerian and Lagrangian derivatives (I)

- The rigorous way to address this problem requires a **notion of differentiation of functions**  $\Omega \mapsto u_\Omega$ , which to a domain  $\Omega$  associate a function defined on  $\Omega$ .
- One could think of two ways of doing so:

### The Eulerian point of view:

For a fixed  $x \in \Omega$ ,  $u'_\Omega(\theta)(x)$  is the derivative of the mapping


$$\theta \mapsto u_{\Omega_\theta}(x).$$



### The Lagrangian point of view:

For a fixed  $x \in \Omega$ ,  $u'_\Omega(\theta)(x)$  is the derivative of the mapping

$$\theta \mapsto u_{\Omega_\theta}((\text{Id} + \theta)(x)).$$



## Eulerian and Lagrangian derivatives (II)

- The Eulerian notion of shape derivative, however more intuitive, is more difficult to define rigorously. In particular, differentiating the **boundary conditions** satisfied by  $u_\Omega$  is awkward:

*Even for “small”  $\theta$ ,  $u_{\Omega_\theta}(x)$  may not make any sense if  $x \in \partial\Omega$ !*

- The Lagrangian derivative  $\dot{u}_\Omega(\theta)$  can be rigorously defined, and lends itself to easier mathematical analysis.
- The rigorous mathematical trail consists in:
  - ① Defining properly the Lagrangian derivative  $\dot{u}_\Omega(\theta)$ ;
  - ② Defining the Eulerian derivative  $u'_\Omega(\theta)$  is **defined** from  $\dot{u}_\Omega(\theta)$ , via the formula:

$$u'_\Omega(\theta) = \dot{u}_\Omega(\theta) - \nabla u_\Omega(x) \cdot \theta,$$

so that the expected **chain rule** holds for the expression  $u_{(\text{Id}+\theta)(\Omega)} \circ (\text{Id} + \theta)$ :

$$\forall x \in \Omega, \dot{u}_\Omega(\theta)(x) = u'_\Omega(\theta)(x) + \nabla u_\Omega(x) \cdot \theta(x).$$

Let  $\Omega \mapsto u_\Omega \in H^1(\Omega)$  be a function which to a domain  $\Omega$ , associates a function  $u_\Omega$  defined on  $\Omega$ .

### Definition 2.

The mapping  $u : \Omega \mapsto u_\Omega$  admits a **material**, or **Lagrangian** derivative  $\dot{u}_\Omega(\theta) \in H^1(\Omega)$  at a particular domain  $\Omega$  provided the **transported function**

$$W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d) \ni \theta \longmapsto \bar{u}(\theta) := u_{\Omega_\theta} \circ (\text{Id} + \theta) \in H^1(\Omega),$$

defined in the neighborhood of  $0 \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$ , is differentiable at  $\theta = 0$ .

This allows to *define* the notion of Eulerian derivative.

### Definition 3.

The mapping  $u : \Omega \mapsto u_\Omega$  has a **Eulerian derivative**  $u'_\Omega(\theta)$  at a given domain  $\Omega$  in the direction  $\theta \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$  if:

- ① It admits a material derivative  $\dot{u}_\Omega(\theta)$  at  $\Omega$ ;
- ② the quantity  $\nabla u_\Omega \cdot \theta$  belongs to  $H^1(\Omega)$ .

One defines then:

$$u'_\Omega(\theta) = \dot{u}_\Omega(\theta) - \nabla u_\Omega \cdot \theta \in H^1(\Omega).$$

## Eulerian and Lagrangian derivatives (V)

Once Lagrangian and Eulerian derivatives are known, the shape derivative of a **quantity of interest** involving  $u_\Omega$  is readily obtained.

### Proposition 6.

Let  $\Omega \subset \mathbb{R}^d$  be a smooth bounded domain, and suppose that  $\Omega \mapsto u_\Omega$  has a **Lagrangian derivative**  $\dot{u}_\Omega$  at  $\Omega$ . If  $j : \mathbb{R} \rightarrow \mathbb{R}$  is regular enough, the function

$$J(\Omega) = \int_{\Omega} j(u_\Omega) \, dx$$

is then **shape differentiable** at  $\Omega$ , and:

$$\forall \theta \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d), \quad J'(\Omega)(\theta) = \int_{\Omega} (j'(u_\Omega)\dot{u}_\Omega(\theta) + (\operatorname{div}\theta)j(u_\Omega)) \, dx.$$

If, in addition,  $\Omega \mapsto u_\Omega$  has a **Eulerian derivative**  $u'_\Omega$  at  $\Omega$ , the **"chain rule"** holds:

$$J'(\Omega)(\theta) = \underbrace{\int_{\partial\Omega} j(u_\Omega) \theta \cdot n \, ds}_{\text{Derivative of the partial mapping } \Omega \mapsto \int_{\Omega} j(u_\Omega)} + \underbrace{\int_{\Omega} j'(u_\Omega) u'_\Omega(\theta) \, dx}_{\text{Derivative of the partial mapping } \Omega \mapsto \int_{\Omega} j(u_\Omega)} .$$



## Eulerian and Lagrangian derivatives (VI)

Idea of the proof: As usual, a **change of variables** yields:

$$J(\Omega_\theta) = \int_{(\text{Id}+\theta)(\Omega)} j(u_{\Omega_\theta}) \, dx = \int_{\Omega} |\det(\text{I} + \nabla\theta)| j(\bar{u}(\theta)) \, dx.$$

- The mapping  $\theta \mapsto |\det(\text{I} + \nabla\theta)|$  is Fréchet differentiable at  $\theta = 0$  and
$$|\det(\text{I} + \nabla\theta)| = 1 + \text{div}\theta + o(\theta);$$

- The mapping  $\theta \mapsto \bar{u}(\theta)$  is Fréchet differentiable at  $\theta = 0$  and
$$\bar{u}(\theta) = u_\Omega + \dot{u}_\Omega(\theta) + o(\theta);$$

Then, using the **chain rule**,  $\theta \mapsto J(\Omega_\theta)$  is Fréchet differentiable at  $\theta = 0$ , and:

$$J'(\Omega)(\theta) = \int_{\Omega} ((\text{div}\theta)j(u_\Omega) + j'(u_\Omega)\dot{u}_\Omega(\theta)) \, dx.$$

Now, if  $\Omega \mapsto u_\Omega$  as a **Eulerian derivative**, the **definition**  $u'_\Omega(\theta) = \dot{u}_\Omega(\theta) - \nabla u_\Omega \cdot \theta$  combined with the Green's formula yields:

$$J'(\Omega)(\theta) = \int_{\partial\Omega} j(u_\Omega) \theta \cdot n \, ds + \int_{\Omega} j'(u_\Omega) u'_\Omega(\theta) \, dx.$$



## Eulerian and Lagrangian derivatives (VII)

The calculation of the shape derivative  $J'(\Omega)(\theta)$  thus rests on those of the Lagrangian and Eulerian derivatives of  $\Omega \mapsto u_\Omega$ , where

$$\begin{cases} -\Delta u_\Omega = f & \text{in } \Omega, \\ u_\Omega = 0 & \text{on } \partial\Omega. \end{cases}$$

The following result characterizes the **Lagrangian derivative** of  $\Omega \mapsto u_\Omega$ .

### Theorem 7.

*The mapping  $\Omega \mapsto u_\Omega \in H_0^1(\Omega)$  has a **Lagrangian derivative**  $\dot{u}_\Omega(\theta)$ , and for any  $\theta \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$ ,  $\dot{u}_\Omega(\theta) \in H_0^1(\Omega)$  is the unique solution to the variational problem:*

$$\begin{aligned} \forall v \in H_0^1(\Omega), \quad \int_{\Omega} \nabla(\dot{u}_\Omega(\theta)) \cdot \nabla v \, dx &= \int_{\Omega} \operatorname{div}(f\theta)v \, dx \\ &\quad - \int_{\Omega} (\operatorname{div}(\theta)\mathbf{I} - \nabla\theta - \nabla\theta^T)\nabla u_\Omega \cdot \nabla v \, dx, \end{aligned}$$

*or, under classical form:*

$$\begin{cases} -\Delta(\dot{u}_\Omega(\theta)) = \operatorname{div}(f\theta) - \operatorname{div}((\operatorname{div}(\theta)\mathbf{I} - \nabla\theta - \nabla\theta^T)\nabla u_\Omega) & \text{in } \Omega, \\ \dot{u}_\Omega(\theta) = 0 & \text{on } \partial\Omega. \end{cases}$$

Idea of the proof:

- The variational problem satisfied by  $u_{\Omega_\theta}$  is:

$$\forall v \in H_0^1(\Omega_\theta), \quad \int_{\Omega_\theta} \nabla u_{\Omega_\theta} \cdot \nabla v \, dx = \int_{\Omega_\theta} f v \, dx.$$

- By a change of variables, the **transported function**  $\bar{u}(\theta) = u_{\Omega_\theta} \circ (\text{Id} + \theta)$  satisfies:

$$\forall v \in H_0^1(\Omega), \quad \int_{\Omega} A(\theta) \nabla \bar{u}(\theta) \cdot \nabla v \, dx = \int_{\Omega} |\det(\text{Id} + \nabla \theta)| (f \circ (\text{Id} + \theta)) v \, dx,$$

where

$$A(\theta) := |\det(\text{Id} + \nabla \theta)| (\text{Id} + \nabla \theta)^{-1} (\text{Id} + \nabla \theta)^{-T}.$$

- This variational problem features a **fixed domain** and a **fixed function space**  $H_0^1(\Omega)$ , and only the **coefficients of the formulation** depend on  $\theta$ .

⇒ This structure lends itself to the use of the strategy based on the **Implicit Function theorem** to calculate the derivative of  $\theta \mapsto \bar{u}(\theta)$ .

## Eulerian and Lagrangian derivatives (IX)

- The problem can now be written as an equation for  $\bar{u}(\theta)$ :

$$\mathcal{F}(\theta, \bar{u}(\theta)) = \mathcal{G}(\theta),$$

for appropriate definitions of the operators:

- $\mathcal{F} : W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d) \times H_0^1(\Omega) \rightarrow H^{-1}(\Omega)$ ,
  - $\mathcal{G} : W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d) \rightarrow H^{-1}(\Omega)$ .
- The **implicit function theorem** shows that  $\theta \mapsto \bar{u}(\theta)$  is differentiable at  $\theta = 0$ .
  - The **Lagrangian derivative**  $\dot{u}_\Omega^\circ(\theta)$  of the **transported mapping**  $\bar{u}(\theta)$  can now be computed by taking derivatives inside the variational formula:

$$\begin{aligned} \forall v \in H_0^1(\Omega), \quad \int_{\Omega} \nabla \dot{u}_\Omega^\circ(\theta) \cdot \nabla v \, dx &= \int_{\Omega} \operatorname{div}(f\theta)v \, dx \\ &\quad - \int_{\Omega} (\operatorname{div}(\theta)\mathbf{I} - \nabla\theta - \nabla\theta^T)\nabla u_\Omega \cdot \nabla v \, dx. \end{aligned}$$



## Eulerian and Lagrangian derivatives (X)

- The **Eulerian derivative** of  $u_\Omega$  can now be computed from its **Lagrangian derivative**. It satisfies (after elementary, but tedious calculations):

$$\begin{cases} -\Delta(u'_\Omega(\theta)) = 0 & \text{in } \Omega, \\ u'_\Omega(\theta) = -(\theta \cdot n) \frac{\partial u_\Omega}{\partial n} & \text{on } \partial\Omega. \end{cases}$$

- At this point, we have thus calculated the **shape derivative** of  $J(\Omega)$  as:

$$J'(\Omega)(\theta) = \int_{\Omega} (j'(u_\Omega) \dot{u}_\Omega(\theta) + (\operatorname{div}\theta)j(u_\Omega)) \, dx,$$

or, involving the Eulerian derivative of  $\Omega \mapsto u_\Omega$ ,

$$J'(\Omega)(\theta) = \int_{\partial\Omega} j(u_\Omega) \theta \cdot n \, ds + \int_{\Omega} j'(u_\Omega) u'_\Omega(\theta) \, dx.$$

- The identification of a **descent direction**  $\theta$  for  $J(\Omega)$  (i.e. such that  $J'(\Omega)(\theta) < 0$ ) is awkward, since  $\dot{u}_\Omega(\theta)$  and  $u'_\Omega(\theta)$  depend **implicitly** on  $\theta$  (via a PDE).

## Eulerian and Lagrangian derivatives (XI): the adjoint method

**Idea:** “Lift” the term of  $J'(\Omega)(\theta)$  which features the Lagrangian (or the Eulerian) derivative of  $u_\Omega$  by introducing an **adequate adjoint problem**.

### Theorem 8.

The shape derivative  $J'(\Omega)(\theta)$  rewrites (**volume form**):

$$J'(\Omega)(\theta) = \int_{\Omega} (\operatorname{div}\theta)j(u_\Omega) \, dx + \int_{\Omega} (\operatorname{div}(\theta)\mathbf{I} - \nabla\theta - \nabla\theta^T)\nabla u_\Omega \cdot \nabla p_\Omega \, dx \\ - \int_{\Omega} \operatorname{div}(f\theta)p_\Omega \, dx,$$

where the **adjoint state**  $p_\Omega \in H_0^1(\Omega)$  is the solution to the equation:

$$\begin{cases} -\Delta p_\Omega = -j'(u_\Omega) & \text{in } \Omega, \\ p_\Omega = 0 & \text{on } \partial\Omega. \end{cases}$$

If  $u_\Omega$  and  $p_\Omega$  are **more regular** ( $u_\Omega, p_\Omega \in H^2(\Omega)$ ), this rewrites under the equivalent **surface form**:

$$J'(\Omega)(\theta) = \int_{\partial\Omega} j(u_\Omega)\theta \cdot n \, ds - \int_{\partial\Omega} \frac{\partial u_\Omega}{\partial n} \frac{\partial p_\Omega}{\partial n} \theta \cdot n \, ds - \int_{\partial\Omega} f p_\Omega \theta \cdot n \, ds.$$

Proof of the volume form.

- The shape derivative  $J'(\Omega)(\theta)$  reads:

$$J'(\Omega)(\theta) = \int_{\Omega} (j'(u_{\Omega}) \dot{u}_{\Omega}(\theta) + (\operatorname{div}\theta)j(u_{\Omega})) \, dx.$$

- Here, the Lagrangian derivative  $\dot{u}_{\Omega}(\theta) \in H_0^1(\Omega)$  solves:

$$\begin{aligned} \forall v \in H_0^1(\Omega), \quad \int_{\Omega} \nabla \dot{u}_{\Omega}(\theta) \cdot \nabla v \, dx &= \int_{\Omega} \operatorname{div}(f\theta)v \, dx \\ &\quad - \int_{\Omega} (\operatorname{div}(\theta)\mathbf{I} - \nabla\theta - \nabla\theta^T)\nabla u_{\Omega} \cdot \nabla v \, dx. \end{aligned}$$

- This is to be compared with the variational formulation for  $p_{\Omega}$ :

$$\forall v \in H_0^1(\Omega), \quad \int_{\Omega} \nabla p_{\Omega} \cdot \nabla v \, dx = - \int_{\Omega} j'(u_{\Omega})v \, dx.$$

- Thus,

$$\begin{aligned} J'(\Omega)(\theta) &= \int_{\Omega} (\operatorname{div}\theta)j(u_{\Omega}) \, dx + \int_{\Omega} j'(u_{\Omega})\dot{u}_{\Omega}(\theta) \, dx, \\ &= \int_{\Omega} (\operatorname{div}\theta)j(u_{\Omega}) \, dx - \int_{\Omega} \nabla p_{\Omega} \cdot \nabla \dot{u}_{\Omega}(\theta) \, dx, \end{aligned}$$

where we have used the variational formulation for  $p_{\Omega}$  with  $\dot{u}_{\Omega}(\theta)$  as test function.

- Now taking  $p_{\Omega}$  as test function in the variational formulation for  $\dot{u}_{\Omega}(\theta)$  yields the desired result:

$$\begin{aligned} J'(\Omega)(\theta) &= \int_{\Omega} (\operatorname{div}\theta)j(u_{\Omega}) \, dx + \int_{\Omega} (\operatorname{div}(\theta)\mathbf{I} - \nabla\theta - \nabla\theta^T)\nabla u_{\Omega} \cdot \nabla p_{\Omega} \, dx \\ &\quad - \int_{\Omega} \operatorname{div}(f\theta)p_{\Omega} \, dx. \end{aligned}$$



Proof of the surface form. The main idea reads as follows:

- Since  $u_\Omega$  and  $p_\Omega \in H^2(\Omega)$ , we perform **integration by parts** in the volume form to end up with an expression of the form:

$$J'(\Omega)(\theta) = \int_{\partial\Omega} v_\Omega \theta \cdot n \, ds + \int_{\partial\Omega} t_\Omega \cdot \theta_{\partial\Omega} \, ds + \int_\Omega S_\Omega \cdot \theta \, dx,$$

where:

- $v_\Omega : \partial\Omega \rightarrow \mathbb{R}$  is a scalar field;
- $t_\Omega : \partial\Omega \rightarrow \mathbb{R}^d$  is a vector field, acting on the **tangential component** of  $\theta$ :

$$\theta_{\partial\Omega} := \theta - (\theta \cdot n)n;$$

- $S_\Omega : \Omega \rightarrow \mathbb{R}^d$  is a vector field,

whose expressions are explicit in terms of  $u_\Omega$  and  $p_\Omega$ .

- If we believe the **Structure theorem**,  $t_\Omega$  and  $S_\Omega$  must equal 0, ... which we verify.
- A tedious calculation eventually yields the result:

$$J'(\Omega)(\theta) = \int_{\partial\Omega} j(u_\Omega) \theta \cdot n \, ds - \int_{\partial\Omega} \frac{\partial u_\Omega}{\partial n} \frac{\partial p_\Omega}{\partial n} \theta \cdot n \, ds - \int_{\partial\Omega} f p_\Omega \theta \cdot n \, ds.$$

## Eulerian and Lagrangian derivatives: volume form vs. surface form

- The volume form is easier to derive, and demands **minimal regularity** from  $u_\Omega$ ,  $p_\Omega$ .
- For this reason, it is often more convenient for studying mathematical properties of shape derivatives (e.g. their finite element approximation).
- The volume form is **explicit in terms of  $\theta$** ... but it does not allow for a straightforward identification of a descent direction.

⇒ Need to rely on the **“Hilbertian trick”** to achieve this.

- The surface form requires **higher regularity** from  $u_\Omega$ ,  $p_\Omega$ , which is often guaranteed by **elliptic regularity**, provided  $\Omega$  and  $f$  are “smooth enough”.
- The surface form has a more compact expression, which explicitly fulfills the **Structure theorem**.

⇒ A descent direction  $\theta$  for  $J(\Omega)$  is immediately revealed.

## Eulerian and Lagrangian derivatives: summary

- Mathematically speaking, the above trail is the **rigorous** way to assess the differentiability of shape functionals.
- As we have seen, the techniques presented above (in particular the adjoint technique) exist in much more general frameworks than shape optimization, and pertain to the framework of **optimal control theory**.
- Calculating shape derivatives by these means requires tedious calculations.
- In practice, a version of **Céa's method** allows for a formal, simpler way to calculate shape derivatives.

## Part III

# Geometric optimization problems

- 1 The method of Hadamard and shape derivatives
- 2 Shape derivatives of PDE-constrained functionals: the rigorous way, using Eulerian and material derivatives
- 3 C ea's method for calculating shape derivatives
- 4 Numerical aspects of geometric methods
- 5 The level set method for shape optimization

As we have seen, the philosophy of **Céa's method** comes from optimization theory:

- We express  $J(\Omega)$  as the value of an  $\Omega$ -dependent Lagrangian functional:

$$\mathcal{L}(\Omega, u, p) = \underbrace{\int_{\Omega} j(u) \, dx}_{\text{Objective function at stake}} + \underbrace{\int_{\Omega} (-\Delta u - f)p \, dx}_{\substack{u=u_{\Omega} \text{ is enforced as a constraint} \\ \text{by penalization with the Lagrange multiplier } p}},$$

at a saddle point  $(u, p) = (u_{\Omega}, p_{\Omega})$ .

- The “parameter”  $\Omega$ , and the variables  $(u, p)$  must be **independent**.
- The nice features of the derivative of a saddle point value with respect to a parameter allow for significant simplifications in the calculation.


This method is **formal**: in particular, it assumes that we already know that  $\Omega \mapsto u_{\Omega}$  is differentiable.

## Céa's method (II)

- The objective function  $J(\Omega)$  is expressed as the value

$$J(\Omega) = \mathcal{L}(\Omega, u_\Omega, p_\Omega),$$

of a suitably defined Lagrangian  $\mathcal{L}(\Omega, u, p)$  at a **saddle point**  $(u_\Omega, p_\Omega)$ .



- The shape derivative  $J'(\Omega)(\theta)$  reads, **formally**:

$$J'(\Omega)(\theta) = \underbrace{\frac{\partial \mathcal{L}}{\partial \Omega}(\Omega, u_\Omega, p_\Omega)(\theta)}_{\substack{\text{Shape derivative of } \Omega \mapsto \mathcal{L}(\Omega, u, p) \\ \text{taken at } (u, p) = (u_\Omega, p_\Omega)}} + \underbrace{\frac{\partial \mathcal{L}}{\partial u}(\Omega, u_\Omega, p_\Omega)(u'_\Omega(\theta))}_{=0} + \underbrace{\frac{\partial \mathcal{L}}{\partial p}(\Omega, u_\Omega, p_\Omega)(p'_\Omega(\theta))}_{=0}.$$

## Céa's method: the Neumann case (I)

We first consider the case of Neumann boundary conditions:

$$\begin{cases} -\Delta u + u = f & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = 0 & \text{on } \partial\Omega, \end{cases}$$

where the  $+u$  term is added for commodity, so that the system is well-posed in  $H^1(\Omega)$  without any further assumption on  $f$ .

Consider the following **Lagrangian functional**:

$$\mathcal{L}(\Omega, u, p) = \underbrace{\int_{\Omega} j(u) \, dx}_{\substack{\text{Objective function} \\ \text{where } u_{\Omega} \text{ is replaced by } u}} + \underbrace{\int_{\Omega} \nabla u \cdot \nabla p \, dx + \int_{\Omega} up \, dx - \int_{\Omega} fp \, dx}_{\substack{\text{Penalization of the "constraint" } u=u_{\Omega}: \\ \int_{\Omega} (-\Delta u + u - f)p \, dx = 0}}$$

which is defined for any shape  $\Omega \in \mathcal{U}_{\text{ad}}$ , and for any  $u, p \in H^1(\mathbb{R}^d)$ , so that the variables  $\Omega$ ,  $u$  and  $p$  are independent.

## Céa's method: the Neumann case (II)

By construction, evaluating  $\mathcal{L}(\Omega, u, p)$  with  $u = u_\Omega$  yields:

$$\forall p \in H^1(\mathbb{R}^d), \mathcal{L}(\Omega, u_\Omega, p) = \int_{\Omega} j(u_\Omega) dx = J(\Omega).$$

For a fixed shape  $\Omega$ , we search for the **saddle points**  $(u, p) \in \mathbb{R}^d \times \mathbb{R}^d$  of  $\mathcal{L}(\Omega, \cdot, \cdot)$ . The first-order necessary conditions read:

- $\forall \hat{p} \in H^1(\mathbb{R}^d), \frac{\partial \mathcal{L}}{\partial p}(\Omega, u, p)(\hat{p}) = \int_{\Omega} \nabla u \cdot \nabla \hat{p} dx + \int_{\Omega} u \hat{p} dx - \int_{\Omega} f \hat{p} dx = 0.$
- $\forall \hat{u} \in H^1(\mathbb{R}^d), \frac{\partial \mathcal{L}}{\partial u}(\Omega, u, p)(\hat{u}) = \int_{\Omega} j'(u) \hat{u} dx + \int_{\Omega} \nabla \hat{u} \cdot \nabla p dx + \int_{\Omega} \hat{u} p dx = 0.$



## Céa's method: the Neumann case (III)

Step 1: Identification of  $u$ :

$$\forall q \in H^1(\mathbb{R}^d), \int_{\Omega} \nabla u \cdot \nabla q \, dx + \int_{\Omega} uq \, dx - \int_{\Omega} fq \, dx = 0.$$

- Taking  $q$  as any  $C^\infty$  function  $\psi$  with compact support in  $\Omega$  yields:

$$\int_{\Omega} \nabla u \cdot \nabla \psi \, dx + \int_{\Omega} u\psi \, dx - \int_{\Omega} f\psi \, dx = 0 \Rightarrow \boxed{-\Delta u + u = f \text{ in } \Omega}.$$

- Now taking  $q$  as any  $C^\infty$  function  $\psi$  and using Green's formula:

$$\int_{\partial\Omega} \frac{\partial u}{\partial n} \psi \, ds = 0 \Rightarrow \boxed{\frac{\partial u}{\partial n} = 0 \text{ on } \partial\Omega}.$$

**Conclusion:**  $u = u_\Omega$ .

**Step 2:** *Identification of  $p$ :*

$$\forall v \in H^1(\mathbb{R}^d), \quad \int_{\Omega} j'(u)v + \int_{\Omega} \nabla v \cdot \nabla p \, dx + \int_{\Omega} vp \, dx = 0.$$

- Taking  $v$  as **any**  $C^\infty$  function  $\psi$  with compact support in  $\Omega$  yields:

$$\int_{\Omega} \nabla p \cdot \nabla \psi \, dx + \int_{\Omega} vp \, dx + \int_{\Omega} j'(u)\psi \, dx = 0 \Rightarrow \boxed{-\Delta p + p = -j'(u_\Omega) \text{ in } \Omega.}$$

- Now taking  $v$  as **any**  $C^\infty$  function  $\psi$  and using Green's formula:

$$\int_{\partial\Omega} \frac{\partial p}{\partial n} \psi \, ds = 0 \Rightarrow \boxed{\frac{\partial p}{\partial n} = 0 \text{ on } \partial\Omega.}$$

**Conclusion:**  $p = p_\Omega$ , solution to  $\begin{cases} -\Delta p + p = -j'(u_\Omega) & \text{in } \Omega, \\ \frac{\partial p}{\partial n} = 0 & \text{on } \partial\Omega. \end{cases}$

Step 3: Calculation of the shape derivative  $J'(\Omega)(\theta)$ :

- We go back to the fact that:

$$\forall q \in H^1(\mathbb{R}^d), \quad \mathcal{L}(\Omega, u_\Omega, q) = \int_{\Omega} j(u_\Omega) \, dx = J(\Omega).$$

- Differentiating with respect to  $\Omega$  yields, thanks to the chain rule:

$$\forall \theta \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d), \quad J'(\Omega)(\theta) = \frac{\partial \mathcal{L}}{\partial \Omega}(\Omega, u_\Omega, q)(\theta) + \frac{\partial \mathcal{L}}{\partial u}(\Omega, u_\Omega, q)(u'_\Omega(\theta)),$$

where  $u'_\Omega(\theta)$  is the **Eulerian derivative** of  $\Omega \mapsto u_\Omega$  (assumed to exist).

- Now, choosing  $q = p_\Omega$  produces, since  $\frac{\partial \mathcal{L}}{\partial u}(\Omega, u_\Omega, p_\Omega) = 0$ :

$$J'(\Omega)(\theta) = \frac{\partial \mathcal{L}}{\partial \Omega}(\Omega, u_\Omega, p_\Omega)(\theta).$$

## Céa's method: the Neumann case (VI)

The last (partial) derivative boils down to that of a functional of the form:

$$\Omega \mapsto \int_{\Omega} f(x) \, dx,$$

where  $f$  is a fixed function.

Using Theorem 2, we end up with:

$$\forall \theta \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d),$$
$$J'(\Omega)(\theta) = \int_{\partial\Omega} \left( j(u_{\Omega}) + \nabla u_{\Omega} \cdot \nabla p_{\Omega} + u_{\Omega} p_{\Omega} - f p_{\Omega} \right) \theta \cdot n \, ds.$$

## Céa's method: the Dirichlet case (I)

- We now consider the problem of calculating the derivative of:

$$J(\Omega) = \int_{\Omega} j(u_{\Omega}) \, dx, \quad \text{where } \begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases} .$$

- **Warning:** When the state  $u_{\Omega}$  satisfies **essential boundary conditions**, i.e. boundary conditions that are tied to the **definition space of functions** (here,  $H_0^1(\Omega)$ ), an additional difficulty generally arises.

- It is no longer possible to rely on the Lagrangian

$$\mathcal{L}(\Omega, u, p) = \int_{\Omega} j(u) \, dx + \int_{\Omega} \nabla u \cdot \nabla p \, dx - \int_{\Omega} fp \, dx,$$

since it would have to be defined for  $u, p \in H_0^1(\Omega)$ .

- In this case, the arguments  $\Omega, u, p$  would not be **independent**.

## Céa's method: the Dirichlet case (II)

**Solution:** Add an extra variable  $\mu \in H^1(\mathbb{R}^d)$  to the Lagrangian to **penalize** the boundary condition: for all  $u, p, \lambda \in H^1(\mathbb{R}^d)$ ;

$$\mathcal{L}(\Omega, u, p, \lambda) = \underbrace{\int_{\Omega} j(u) \, dx}_{\text{Objective function where } u_{\Omega} \text{ is replaced by } u} + \underbrace{\int_{\Omega} (-\Delta u - f)p \, dx}_{\text{penalization of the "constraint" } -\Delta u = f} + \underbrace{\int_{\partial\Omega} \lambda u \, ds}_{\text{penalization of the "constraint" } u=0 \text{ on } \partial\Omega} .$$

By Green's formula,  $\mathcal{L}(u, p, \lambda)$  rewrites:

$$\mathcal{L}(\Omega, u, p, \lambda) = \int_{\Omega} j(u) \, dx + \int_{\Omega} \nabla u \cdot \nabla p \, dx - \int_{\Omega} fp \, dx + \int_{\partial\Omega} \left( \lambda u - \frac{\partial u}{\partial n} p \right) \, ds.$$

Of course, evaluating  $\mathcal{L}(u, p, \lambda)$  with  $u = u_{\Omega}$ , it comes:

$$\forall p, \lambda \in H^1(\mathbb{R}^d), \quad \mathcal{L}(\Omega, u_{\Omega}, p, \lambda) = \int_{\Omega} j(u_{\Omega}) \, dx.$$

## Céa's method: the Dirichlet case (III)

For a fixed shape  $\Omega$ , we look for the **saddle points**  $(u, p, \lambda) \in (H^1(\mathbb{R}^d))^3$  of the functional  $\mathcal{L}(\Omega, \cdot, \cdot, \cdot)$ . The first-order necessary conditions are:

- $\forall \hat{p} \in H^1(\mathbb{R}^d)$ ,  $\frac{\partial \mathcal{L}}{\partial p}(\Omega, u, p, \lambda)(\hat{p}) = \int_{\Omega} \nabla u \cdot \nabla \hat{p} \, dx - \int_{\Omega} f \hat{p} \, dx + \int_{\partial\Omega} \frac{\partial u}{\partial n} \hat{p} \, ds = 0$ .
- $\forall \hat{u} \in H^1(\mathbb{R}^d)$ ,  $\frac{\partial \mathcal{L}}{\partial u}(\Omega, u, p, \lambda)(\hat{u}) = \int_{\Omega} j'(u) \hat{u} \, dx + \int_{\Omega} \nabla \hat{u} \cdot \nabla p \, dx + \int_{\partial\Omega} \left( \lambda \hat{u} - \frac{\partial \hat{u}}{\partial n} p \right) \, ds = 0$ .
- $\forall \hat{\lambda} \in H^1(\mathbb{R}^d)$ ,  $\frac{\partial \mathcal{L}}{\partial \lambda}(\Omega, u, p, \lambda)(\hat{\lambda}) = \int_{\partial\Omega} \hat{\lambda} u \, ds = 0$ .

Step 1: Identification of  $u$ :

$$\forall q \in H^1(\mathbb{R}^d), \int_{\Omega} \nabla u \cdot \nabla q \, dx - \int_{\Omega} f q \, dx + \int_{\partial\Omega} \frac{\partial u}{\partial n} q \, ds = 0.$$

- Taking  $q$  as any  $C^\infty$  function  $\psi$  with compact support in  $\Omega$  yields:

$$\forall \psi \in C_c^\infty(\Omega), \int_{\Omega} \nabla u \cdot \nabla \psi \, dx = \int_{\Omega} f \psi \, dx \Rightarrow \boxed{-\Delta u = f \text{ in } \Omega.}$$

- Using  $\frac{\partial \mathcal{L}}{\partial \lambda}(\Omega, u, p, \lambda)(\mu) = 0$  for any  $\mu = \psi \in C_c^\infty(\mathbb{R}^d)$  yields:

$$\forall \psi \in C_c^\infty(\mathbb{R}^d), \int_{\partial\Omega} \psi u \, ds = 0 \Rightarrow \boxed{u = 0 \text{ on } \partial\Omega.}$$

**Conclusion:**  $u = u_\Omega$ .



## Step 2: Identification of $p$ :

$$\forall v \in H^1(\mathbb{R}^d), \int_{\Omega} j'(u)v \, dx + \int_{\Omega} \nabla v \cdot \nabla p \, dx + \int_{\partial\Omega} \left( \lambda v - \frac{\partial v}{\partial n} p \right) \, ds = 0.$$

- Taking  $q$  as any  $C^\infty$  function  $\psi$  with compact support in  $\Omega$  yields:

$$\forall \psi \in C_c^\infty(\Omega), \int_{\Omega} \nabla p \cdot \nabla \psi \, dx + \int_{\Omega} j'(u)\psi \, dx = 0$$

$$\Rightarrow \boxed{-\Delta p = -j'(u_\Omega) \text{ in } \Omega.}$$

- Now taking  $v$  as a  $C^\infty$  function  $\psi$  and using Green's formula:

$$\forall \psi \in C_c^\infty(\mathbb{R}^d), \int_{\partial\Omega} \frac{\partial p}{\partial n} \psi \, ds + \int_{\partial\Omega} \left( \lambda \psi - \frac{\partial \psi}{\partial n} p \right) \, ds = 0.$$

Step 2 (continued):

- Varying the normal trace  $\frac{\partial \psi}{\partial n}$  while imposing  $\psi = 0$  on  $\partial\Omega$ , one gets:

$$p = 0 \text{ on } \partial\Omega.$$

**Conclusion:**  $p = p_\Omega$ , solution to 
$$\begin{cases} -\Delta p = -j'(u_\Omega) & \text{in } \Omega, \\ p = 0 & \text{on } \partial\Omega. \end{cases}$$

- In addition, varying the trace of  $\psi$  on  $\partial\Omega$  while imposing  $\frac{\partial \psi}{\partial n} = 0$ :

$$\lambda_\Omega = -\frac{\partial p_\Omega}{\partial n} \text{ on } \partial\Omega.$$

Step 3: Calculation of the shape derivative  $J'(\Omega)(\theta)$ :

- We return to the fact that:

$$\forall q, \mu \in H^1(\mathbb{R}^d), \quad \mathcal{L}(\Omega, u_\Omega, q, \mu) = \int_{\Omega} j(u_\Omega) \, dx.$$

- Differentiating with respect to  $\Omega$  yields, for all  $\theta \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$ :

$$J'(\Omega)(\theta) = \frac{\partial \mathcal{L}}{\partial \Omega}(\Omega, u_\Omega, q, \mu)(\theta) + \frac{\partial \mathcal{L}}{\partial u}(\Omega, u_\Omega, q, \mu)(u'_\Omega(\theta)),$$

where  $u'_\Omega(\theta)$  is the **Eulerian derivative** of  $\Omega \mapsto u_\Omega$ .

- Taking  $q = p_\Omega$ ,  $\mu = \lambda_\Omega$  produces, since  $\frac{\partial \mathcal{L}}{\partial u}(\Omega, u_\Omega, p_\Omega, \lambda_\Omega) = 0$ :

$$J'(\Omega)(\theta) = \frac{\partial \mathcal{L}}{\partial \Omega}(\Omega, u_\Omega, p_\Omega, \lambda_\Omega)(\theta).$$

## Céa's method: the Dirichlet case (VIII)

Again, this (partial) shape derivative combines derivatives of functions of the form:

$$\Omega \mapsto \int_{\Omega} f(x) \, dx, \text{ or } \Omega \mapsto \int_{\partial\Omega} g(x) \, ds,$$

where  $f$  and  $g$  are fixed functions.

Using Theorems 2 and 3 (and after some calculation), we end up with:

$$\forall \theta \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d), \quad J'(\Omega)(\theta) = \int_{\partial\Omega} \left( j(u_{\Omega}) - \frac{\partial u_{\Omega}}{\partial n} \frac{\partial p_{\Omega}}{\partial n} \right) \theta \cdot n \, ds.$$

## Part III

# Geometric optimization problems

- 1 The method of Hadamard and shape derivatives
- 2 Shape derivatives of PDE-constrained functionals: the rigorous way, using Eulerian and material derivatives
- 3 C ea's method for calculating shape derivatives
- 4 Numerical aspects of geometric methods**
- 5 The level set method for shape optimization

## The generic numerical algorithm

**Initialization:** Start from an initial shape  $\Omega^0$ .

**For**  $n = 0, \dots$  **convergence,**

- 1 Calculate the **state**  $u_{\Omega^n}$  (and the **adjoint**  $p_{\Omega^n}$  if need be) on  $\Omega^n$ .
- 2 Compute the shape derivative  $J'(\Omega^n)$  by evaluating the mathematical formula, and infer a **descent direction**  $\theta^n$  for  $J(\Omega)$ .
- 3 **Advect** the shape  $\Omega^n$  along the displacement field  $\theta^n$ , for a small **pseudo-time step**  $\tau^n$ , so as to obtain

$$\Omega^{n+1} = (\text{Id} + \tau^n \theta^n)(\Omega^n).$$


## One possible implementation

- Each shape  $\Omega^n$  is represented by a **simplicial mesh**  $\mathcal{T}^n$  (i.e. composed of triangles in  $2d$  and of tetrahedra in  $3d$ ).
- The **Finite Element method** is used on  $\mathcal{T}^n$  for computing  $u_{\Omega^n}$  (and  $p_{\Omega^n}$ ).
- The descent direction  $\theta^n$  is obtained from the **surface form** of the shape derivative:

$$J'(\Omega)(\theta) = \int_{\partial\Omega} v_{\Omega}\theta \cdot n \, ds \quad \Rightarrow \quad \theta^n = -v_{\Omega}n \text{ on } \partial\Omega.$$

- The **shape advection** step  $\Omega^n \xrightarrow{\text{Id} + \tau^n \theta^n} \Omega^{n+1}$  is performed by **pushing the nodes** of  $\mathcal{T}^n$  along  $\tau^n \theta^n$ , to obtain the new mesh  $\mathcal{T}^{n+1}$ :

$$\forall \text{ vertex } x \in \mathcal{T}^n, \quad x \mapsto x + \tau^n \theta^n(x).$$



*Deformation of a mesh by relocating its nodes to a prescribed final position.*

## Numerical examples (I)

- In the context of **linear elasticity**, one aims at minimizing the **compliance**  $C(\Omega)$  of a cantilever beam:

$$C(\Omega) = \int_{\Omega} A e(u_{\Omega}) : e(u_{\Omega}) dx.$$

- An equality constraint on the **volume**  $\text{Vol}(\Omega)$  of shapes is imposed by means of a **fixed penalization** procedure.



## Numerical examples (II)

- In the context of **fluid mechanics** (Stokes equations), one aims at minimizing the **viscous dissipation**  $D(\Omega)$  in a pipe:

$$D(\Omega) = 2\nu \int_{\Omega} D(u_{\Omega}) : D(u_{\Omega}) \, dx.$$


- A **volume constraint** is imposed by a *fixed* penalization of the function  $D(\Omega)$ .

## Numerical examples (III)

- Still in **fluid mechanics**, the **viscous dissipation**  $D(\Omega)$  of a double pipe system is minimized.
- A **volume constraint** is imposed.

### I - Existence of many local minimizers:


- In “most” shape optimization problems, no “true” global minimizer exists: the latter would have to be searched as a **homogenized design**;
- However, there exist many **local minimizers**;
- In practice, shape optimization algorithms are **very sensitive** to the initial design, to the size of the computational mesh, etc.



*Several optimized cantilever beams associated to different initial designs.*

### // - The difficulty of mesh deformation:

- The **update of the shape** at each step  $\Omega^n \mapsto (\text{Id} + \theta^n)(\Omega^n) = \Omega^{n+1}$  is realized by **relocating each node**  $x \in \mathcal{T}^n$  to  $x + \tau^n \theta^n(x) \in \mathcal{T}^{n+1}$ .
- This may prove difficult, partly because it may cause **inversion of elements**, resulting in an **invalid** mesh.



*Pushing nodes according to  $\theta^n$  may result in an invalid configuration.*

- For this reason, mesh deformation methods are generally preferred for accounting for “**small displacements**”.

### III - Velocity extension:

- A descent direction  $\theta = -v_\Omega n$  from a shape  $\Omega$  is inferred from the formula:

$$J'(\Omega)(\theta) = \int_{\partial\Omega} v_\Omega(\theta \cdot n) \, ds.$$


- The new shape  $(\text{Id} + \theta)(\Omega)$  only depends on these values of  $\theta$  on  $\partial\Omega$ .
- For many reasons, in numerical practice, it is crucial to **extend  $\theta$**  to  $\Omega$  (or even  $\mathbb{R}^d$ ) in a “clever” way.

*(for instance, deforming a mesh of  $\Omega$  using a “nice” vector field  $\theta$  defined on the whole  $\Omega$  may considerably ease the process)*

- The “natural” extension of the formula  $\theta = -v_\Omega n$ , **which is only legitimate on  $\partial\Omega$**  may not be a “good” choice.

### IV - Velocity regularization:

- The descent direction  $\theta = -v_{\Omega}n$  on  $\partial\Omega$  may be very **irregular**, because of
  - **numerical artifacts** arising during the finite element analyses.
  - an inherent lack of regularity of  $J'(\Omega)$  for the problem at stake.
- In numerical practice, it is often necessary to **smooth** this descent direction so that the considered shapes stay regular.



*Irregularity of the shape derivative in the very sensitive problem of drag minimization of an airfoil (Taken from [MoPir]). In one iteration, using the unsmoothed shape derivative of  $J(\Omega)$  produces large undesirable artifacts.*

### A popular idea: extend AND regularize the velocity field

- Suppose we aim at extending the *scalar* field  $v_\Omega : \partial\Omega \rightarrow \mathbb{R}$  to  $\Omega$ .
- **Idea:** ( $\approx$  Laplacian smoothing) Trade the “natural” inner product over  $L^2(\partial\Omega)$  for a **more regular** inner product over functions on  $\Omega$ .
- **Example:** Search the extended / regularized scalar field  $V$  as:  
Find  $V \in H^1(\Omega)$  s.t.  $\forall w \in H^1(\Omega)$ ,

$$\alpha^2 \int_{\Omega} \nabla V \cdot \nabla w \, dx + \int_{\Omega} V w \, dx = \int_{\partial\Omega} v_\Omega w \, ds.$$

- The **regularizing parameter**  $\alpha$  controls the balance between the fidelity of  $V$  to  $v_\Omega$  and the intensity of smoothing.

## The Hilbertian method (II)

- The resulting scalar field  $V$  is inherently defined on  $\Omega$  and more regular than  $v_\Omega$ .
- Multiple other **regularizing problems** are possible, associated to different inner products or different function spaces.
- A similar process also allows to:
  - extend  $v_\Omega$  to a large computational box  $D$  (an inner product over functions defined on  $D$  is used),
  - extend the **vector velocity**  $\theta = -v_\Omega n$  to  $\Omega$  or  $D$  (an inner product over vector functions is used, e.g. that of linear elasticity).
- In particular, such a procedure allows to obtain a descent direction from the **volume form** of the shape derivative:

$$J'(\Omega)(\theta) = \int_{\Omega} (r_\Omega \cdot \theta + S_\Omega : \nabla \theta) \, dx,$$

where the fields  $r_\Omega : \Omega \rightarrow \mathbb{R}^d$ ,  $S_\Omega : \Omega \rightarrow \mathbb{R}^{d \times d}$  are known.



## Part III

# Geometric optimization problems


- 1 The method of Hadamard and shape derivatives
- 2 Shape derivatives of PDE-constrained functionals: the rigorous way, using Eulerian and material derivatives
- 3 C ea's method for calculating shape derivatives
- 4 Numerical aspects of geometric methods
- 5 The level set method for shape optimization

## The level set method

**A paradigm:** *the motion of an evolving domain is conveniently described in an **implicit** way.*

A domain  $\Omega \subset \mathbb{R}^d$  is equivalently defined by a function  $\phi : \mathbb{R}^d \rightarrow \mathbb{R}$  such that:

$$\phi(x) < 0 \quad \text{if } x \in \Omega \quad ; \quad \phi(x) = 0 \quad \text{if } x \in \partial\Omega \quad ; \quad \phi(x) > 0 \quad \text{if } x \in \mathring{\Omega}$$




A domain  $\Omega \subset \mathbb{R}^2$  (left), some level sets of an associated level set function (right).

## Level set functions and geometry (I)

If  $\phi : \mathbb{R}^d \rightarrow \mathbb{R}$  is a **level set function** of class  $\mathcal{C}^2$  for  $\Omega$ , such that  $\nabla\phi(x) \neq 0$  on a neighborhood of  $\partial\Omega$ ,

- The **normal vector**  $n$  to  $\partial\Omega$  pointing outward  $\Omega$  reads:

$$\forall x \in \partial\Omega, \quad n(x) = \frac{\nabla\phi(x)}{|\nabla\phi(x)|}.$$



*Normal vector to a domain  $\Omega$ ; some isolines of the function  $\phi$  are dotted.*


## Level set functions and geometry (II)

- The **second fundamental form**  $\mathbb{II}$  of  $\partial\Omega$  is:

$$\forall x \in \partial\Omega, \mathbb{II}(x) = \nabla \left( \frac{\nabla\phi(x)}{|\nabla\phi(x)|} \right).$$

- The **mean curvature**  $\kappa$  of  $\partial\Omega$  is:

$$\forall x \in \partial\Omega, \kappa(x) = \operatorname{div} \left( \frac{\nabla\phi(x)}{|\nabla\phi(x)|} \right).$$



$\mathbb{II}_x(v, v)$  is the curvature of a curve drawn on  $\partial\Omega$  with tangent vector  $v$  at  $x$ .

## Evolution of domains with the level set method

- Let  $\Omega(t) \subset \mathbb{R}^d$  be a domain moving along a velocity field  $v(t, x) \in \mathbb{R}^d$ .
- Let  $\phi(t, x)$  be a level set function for  $\Omega(t)$ .
- The motion of  $\Omega(t)$  translates in terms of  $\phi$  as the **level set advection equation**:


$$\frac{\partial \phi}{\partial t}(t, x) + v(t, x) \cdot \nabla \phi(t, x) = 0$$

- If  $v(t, x)$  is normal to the boundary  $\partial\Omega(t)$ , i.e.:

$$v(t, x) := V(t, x) \frac{\nabla \phi(t, x)}{|\nabla \phi(t, x)|},$$

this rewrites as a **Hamilton-Jacobi equation**:

$$\frac{\partial \phi}{\partial t}(t, x) + V(t, x) |\nabla \phi(t, x)| = 0$$



## The level set method in the context of shape optimization (I)

- A **fixed** computational domain  $D$  is meshed once and for all (e.g. with triangular or quadrilateral elements).
- Each shape  $\Omega^n$  is represented by a **level set function**  $\phi^n$ , defined at the nodes of the mesh.
- As soon as a descent direction  $\theta^n$  from  $\Omega^n$  is available, the **advection step**


$$\Omega^n \mapsto \Omega^{n+1} = (\text{Id} + \tau^n \theta^n)(\Omega^n)$$

is achieved by solving the **advection-like equation**

$$\begin{cases} \frac{\partial \phi}{\partial t} + \theta^n \cdot \nabla \phi = 0 & t \in (0, \tau^n), x \in D \\ \phi(0, \cdot) = \phi^n \end{cases}$$

or if  $\theta^n = v^n n$  is normal, the **Hamilton-Jacobi equation**:

$$\begin{cases} \frac{\partial \phi}{\partial t} + v^n |\nabla \phi| = 0 & t \in (0, \tau^n), x \in D \\ \phi(0, \cdot) = \phi^n \end{cases}$$



Shape accounted for by a level set description (from [AlJouToa])

## The level set method in the context of shape optimization (II)

**Problem:** At each iteration  $n$ , **no mesh** of  $\Omega^n$  is available to solve the finite element problems needed in the calculation of the shape gradient.

**Solution:** The state and adjoint PDE problems posed on  $\Omega^n$  are **approximated** by a problem posed on the whole box  $D$

⇒ Use of a **Fictitious domain method**.

## The ersatz material approximation in linearized elasticity (I)

- In the linear elasticity context, the optimized part of the boundary  $\Gamma$  (i.e. that represented with the level set method) is often **traction-free**.
- The **ersatz material method** approximates the elastic displacement  $u_\Omega : \Omega \rightarrow \mathbb{R}^d$  by that  $u_{\Omega,\varepsilon} : D \rightarrow \mathbb{R}^d$  of the total domain  $D$  when the void  $D \setminus \bar{\Omega}$  is filled with a very 'soft' material:

$$\left\{ \begin{array}{ll} -\operatorname{div}(Ae(u_\Omega)) = 0 & \text{in } \Omega, \\ u_\Omega = 0 & \text{on } \Gamma_D, \\ Ae(u_\Omega)n = g & \text{on } \Gamma_N, \\ Ae(u_\Omega)n = 0 & \text{on } \Gamma. \end{array} \right. \approx \left\{ \begin{array}{ll} -\operatorname{div}(A_\varepsilon e(u_{\Omega,\varepsilon})) = 0 & \text{in } D, \\ u_{\Omega,\varepsilon} = 0 & \text{on } \Gamma_D, \\ A_\varepsilon e(u_{\Omega,\varepsilon})n = g & \text{on } \Gamma_N, \\ Ae(u_{\Omega,\varepsilon})n = 0 & \text{on } \partial D \setminus (\Gamma_D \cup \Gamma_N), \end{array} \right.$$


(Problem posed on  $\Omega$ ) (Problem posed on  $D$ )

where the **approximate Hooke's tensor**  $A_\varepsilon$  reads:


$$A_\varepsilon = \chi_\Omega A + (1 - \chi_\Omega)\varepsilon A, \quad \varepsilon \ll 1.$$





# The ersatz material approximation in linearized elasticity (II)




*Physical situation of a bridge*



*Deformed configuration of the bridge*



*Implicit definition of the bridge on a mesh of  $D$*



*Deformed configuration of the domain  $D$*

## Example: optimization of a 2d bridge using the level set method

- In the context of **linear elasticity**, the **compliance** of a bridge is minimized

$$C(\Omega) = \int_{\Omega} A e(u_{\Omega}) : e(u_{\Omega}) \, dx.$$

- A constraint on the **volume**  $\text{Vol}(\Omega)$  of shapes is imposed.

# Technical appendix

# The Sobolev space $W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$

## Definition 4.

The space  $W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$  is equivalently defined as:

- The space of **bounded** and **Lipschitz** vector fields  $\theta : \mathbb{R}^d \rightarrow \mathbb{R}^d$ , i.e. there exists  $C > 0$  such that:

$$\forall x \in \mathbb{R}^d, |\theta(x)| \leq C, \text{ and } \forall x, y \in \mathbb{R}^d, |\theta(x) - \theta(y)| \leq C|x - y|.$$

- The **Sobolev space** of uniformly bounded functions, with uniformly bounded derivatives:

$$\left\{ \theta \in L^\infty(\mathbb{R}^d), \frac{\partial \theta_i}{\partial x_j} \in L^\infty(\mathbb{R}^d), i, j = 1, \dots, d \right\}.$$

The space  $W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$  is equipped with the norm:

$$\begin{aligned} \|\theta\|_{W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)} &= \sup_{\substack{x, y \in \mathbb{R}^d \\ x \neq y}} \left( |\theta(x)| + \frac{|\theta(x) - \theta(y)|}{|x - y|} \right) \\ &= \|\theta\|_{L^\infty(\mathbb{R}^d)} + \sup_{x \in \mathbb{R}^d} |\nabla \theta(x)|. \end{aligned}$$

## Change of variable formulas (I)

The next theorem is an extension of the usual **change of variables** formula (involving a  $C^1$  diffeomorphism) to the case of a **Lipschitz** diffeomorphism; see [EGar], Chap. 3.

### Theorem 9 (Lipschitz change of variables in volume integrals).

Let  $\Omega \subset \mathbb{R}^d$  be a Lipschitz bounded domain, and  $\varphi : \Omega \rightarrow \mathbb{R}^d$  be a Lipschitz diffeomorphism of  $\mathbb{R}^d$ . Then, for any function  $f \in L^1(\varphi(\Omega))$ ,  $f \circ \varphi$  is in  $L^1(\Omega)$  and:

$$\int_{\varphi(\Omega)} f \, dx = \int_{\Omega} |\det(\nabla \varphi)| f \circ \varphi \, dx.$$

**Remark:** The Jacobian determinant  $|\det(\nabla \varphi)|$  exists a.e. in  $\Omega$ , as a consequence of the **Rademacher theorem**:

*A Lipschitz function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is almost everywhere differentiable.*

## Change of variable formulas (II)

The following theorem is a version of the change of variables formula adapted to [surface integrals](#); see [HenPi], Prop. 5.4.3.

### Theorem 10 (Change of variables in surface integrals).

Let  $\Omega \subset \mathbb{R}^d$  be a bounded domain of class  $C^1$  with boundary  $\Gamma$  and unit normal vector  $n$  pointing outward  $\Omega$ . Let  $\varphi : \Omega \rightarrow \mathbb{R}^d$  be a  $C^1$  diffeomorphism of  $\mathbb{R}^d$ . Then, for any function  $g \in L^1(\varphi(\Gamma))$ ,  $g \circ \varphi$  belongs to  $L^1(\Gamma)$  and:

$$\int_{\varphi(\Gamma)} g \, ds = \int_{\Gamma} |\text{Com}(\nabla\varphi)n| g \circ \varphi \, ds,$$

where  $\text{Com}(M)$  is the cofactor matrix of a  $d \times d$  matrix.

**Remark:** The integrand


$$|\text{Com}(\nabla\varphi)n| = |\det(\nabla\varphi)| |\nabla\varphi^{-T}n|$$

is sometimes called the **tangential Jacobian** of the diffeomorphism  $\varphi$ .

## Surfaces and curvature (I)


At first order, in the neighborhood of a point  $p \in \Gamma$ , a surface  $\Gamma$  behaves like a plane, the **tangent plane**,

- With **normal vector**  $n(p)$ ,
- Which contains the **tangential directions** to  $\Gamma$ .



## Surfaces and curvature (II)

- At second order in the neighborhood of  $p \in \Gamma$ , the surface  $\Gamma$  has one **curvature** in each tangential direction.
- The **principal directions** at  $p$  are those tangential directions  $v_1(p)$  et  $v_2(p)$  associated to the lower and larger curvatures  $\kappa_1(p)$  et  $\kappa_2(p)$ .
- The **mean curvature**  $\kappa(p)$  is the sum  $\kappa(p) = \kappa_1(p) + \kappa_2(p)$ .





## The implicit function theorem

Let us recall the **implicit function theorem**; see [La], Chap. I, Th. 5.9.

### Theorem 11 (Implicit function theorem).

Let  $\Theta, E, F$  be Banach spaces,  $\mathcal{V} \subset \Theta$ ,  $U \subset E$  be open sets. and  $\mathcal{F} : \mathcal{V} \times U \rightarrow G$  be a function of class  $C^p$  for  $p \geq 1$ . Let  $(\theta_0, u_0) \in \mathcal{V} \times U$  be such that  $\mathcal{F}(\theta_0, u_0) = 0$  and assume that:

$d_u \mathcal{F}(\theta_0, u_0) : F \rightarrow G$  is a linear isomorphism.

Then there exist an open subset  $\mathcal{V}' \subset \mathcal{V}$  of  $\theta_0$  in  $\Theta$  and a mapping  $g : \mathcal{V}' \rightarrow U$  of class  $C^p$  satisfying the properties:

- 1  $g(\theta_0) = u_0$ ,
- 2 For all  $\theta \in \mathcal{V}'$ , the equation  $\mathcal{F}(\theta, u) = 0$  has a unique solution  $u \in E$ , given by  $u = g(\theta)$ .

## A glimpse of elliptic regularity

- Existence and uniqueness of the solution  $u$  to an **elliptic equation** (e.g. the **conductivity equation**, the **linear elasticity system**) is often guaranteed by the **Lax-Milgram theory**.
- In general, this theory only supplies “weak” solutions, in a Sobolev space with “low” regularity (typically  $H^1(\Omega)$ ).
- It turns out that this solution is in general “as regular as permitted by the data”.
- **Elliptic regularity** is a general phenomenon, which roughly states:

The solution  $u$  to a second-order **elliptic equation** posed in a **smooth** domain  $\Omega$ , with **smooth** coefficients, is twice more regular than the data  $f$ :

$$f \in H^k(\Omega) \Rightarrow u \in H^{k+2}(\Omega), \text{ and } \|u\|_{H^{k+2}(\Omega)} \leq C\|f\|_{H^k(\Omega)}.$$

## Theorem 12.

Let  $\Omega \subset \mathbb{R}^d$  be a bounded domain of class  $C^{k+2}$ , and let  $f \in H^k(\Omega)$ . Then, the unique solution  $u \in H_0^1(\Omega)$  to the equation

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

belongs to  $H^{k+2}(\Omega)$ , and the following estimate holds:

$$\|u\|_{H^{k+2}(\Omega)} \leq C \|f\|_{H^k(\Omega)},$$

for a constant  $C > 0$  which only depends on  $k$  and  $\Omega$ .

- This is an avatar of a very general phenomenon; similar statements hold for
  - Other types of **boundary conditions** (Neumann, Robin, ...),
  - Other **equations**: the **linearized elasticity system**, the **Stokes equations**, etc.
- We only provide a short sketch of proof; see [?], §9.6 for a comprehensive treatment.

## Sketch of proof

Hint of proof: We proceed in three steps:

- (i) Interior regularity: We prove that for every cut-off function  $\chi \in C_c^\infty(\Omega)$ ,

$$\chi u \in H^2(\Omega), \text{ and } \|\chi u\|_{H^2(\Omega)} \leq C \|f\|_{L^2(\Omega)},$$

for a constant  $C > 0$  depending only on  $\chi$  and  $\Omega$ .

- (ii) Regularity near the boundary: We prove that for any point  $x_0 \in \partial\Omega$ , there exists a bounded open set  $\mathcal{O}$  containing  $x_0$  such that for any cutoff function  $\chi \in C_c^\infty(\mathbb{R}^d)$  with compact support inside  $\overline{\mathcal{O}}$ ,

$$\chi u \in H^2(\Omega), \text{ and } \|\chi u\|_{H^2(\Omega)} \leq C \|f\|_{L^2(\Omega)}.$$

- (iii) Global regularity: Using a **partition of unity** argument, we “glue” the local results from Steps (i) and (ii).

## Proof of the interior regularity statement (I)

### Proof of Step (i): Interior regularity

- By a simple calculation, the function  $\chi u$  satisfies the equation:

$$-\Delta(\chi u) = g, \text{ where } g := -(\Delta\chi)u - 2\nabla\chi \cdot \nabla u - \chi f \in L^2(\mathbb{R}^d). \quad (\text{SF})$$

Under variational form,  $\chi u$  is the unique solution in  $H_0^1(\Omega)$  to the problem:

$$\forall v \in H_0^1(\Omega), \int_{\Omega} \nabla(\chi u) \cdot \nabla v \, dx = \int_{\Omega} g v \, dx. \quad (\text{VF})$$

- **Intuitively**, because  $g \in L^2(\Omega)$  and  $\text{supp}(g)$  is a compact of  $\Omega$ , for  $i = 1, \dots, d$ ,  $\frac{\partial g}{\partial x_i} \in H^{-1}(\Omega)$ . By the standard **Lax-Milgram theory**, the variational problem

$$\forall v \in H_0^1(\Omega), \int_{\Omega} \nabla w_i \cdot \nabla v \, dx = \left\langle \frac{\partial g}{\partial x_i}, v \right\rangle_{H^{-1}(\Omega), H_0^1(\Omega)},$$

obtained by formally taking derivatives in (SF) or (VF), has a unique solution  $w_i \in H_0^1(\Omega)$ , which it is tempting to identify with  $\frac{\partial}{\partial x_i}(\chi u)$ .

- Making this argument **rigorous** relies on the **method of translations** of L. Nirenberg.

## Proof of the interior regularity statement (II)

For a function  $u : \Omega \rightarrow \mathbb{R}$ , a point  $x \in \Omega$ , and a direction  $h \in \mathbb{R}^d$  such that  $|h| < d(x, \partial\Omega)$ , we define the **difference quotient**:

$$D_h u(x) = \frac{u(x+h) - u(x)}{|h|}.$$

### Theorem 13 (The method of translations).

The following statements are equivalent:

- 1  $u \in H^1(\Omega)$ ;
- 2 There exists  $C > 0$  such that:

$$\forall i = 1, \dots, d, \quad \forall \varphi \in C_c^\infty(\Omega), \quad \left| \int_{\Omega} u \frac{\partial \varphi}{\partial x_i} dx \right| \leq C \|\varphi\|_{L^2(\Omega)}.$$

- 3 There exists  $C > 0$  such that for any open subset  $\omega \Subset \Omega$ , and any vector  $h \in \mathbb{R}^d$  with  $|h| < \text{dist}(\omega, \partial\Omega)$ ,

$$\|D_h u\|_{L^2(\omega)} \leq C.$$

In addition, one may take  $C = \|\nabla u\|_{L^2(\Omega)^2}$  in the last two statements.

## Proof of the interior regularity statement (III)

- Taking  $v = D_{-h}D_h(\chi u)$  as test function in the variational formulation for  $\chi u$  is possible because  $\text{supp}(\chi u)$  is a compact of  $\Omega$ ; this yields:

$$\int_{\Omega} \nabla(\chi u) \cdot \nabla(D_{-h}D_h(\chi u)) \, dx = \int_{\Omega} g D_{-h}D_h(\chi u) \, dx.$$

- Performing a discrete integration by parts (i.e. a change of variables), we get:

$$\int_{\Omega} \nabla(D_h(\chi u)) \cdot \nabla(D_h(\chi u)) \, dx = \int_{\Omega} g D_{-h}D_h(\chi u) \, dx.$$

The Cauchy-Schwarz inequality and the translation theorem ((i)  $\Rightarrow$  (iii)) lead to:

$$\|\nabla(D_h(\chi u))\|_{L^2(\Omega)^2}^2 \leq \|g\|_{L^2(\Omega)} \|\nabla(D_h(\chi u))\|_{L^2(\Omega)^2},$$

and so:


$$\|D_h(\nabla(\chi u))\|_{L^2(\Omega)^2} \leq \|g\|_{L^2(\Omega)}.$$

- Eventually, the translation theorem ((iii)  $\Rightarrow$  (i)) implies from this inequality that  $\nabla(\chi u) \in H^1(\Omega)^d$  with the desired estimate.

# Proof of the boundary regularity statement

## Proof of Step (ii):

- Let  $x_0 \in \partial\Omega$ . Because  $\partial\Omega$  is “smooth”, we may take  $\mathcal{O}$  so small that  $\partial\Omega$  is “nearly flat” around  $x_0$  (say,  $\Omega$  coincide with the lower half-space near  $x_0$ ).



- The same argument as before (with “horizontal” translations  $h$ ), shows that:

$$\forall i = 1, \dots, d-1, \quad \frac{\partial(\chi u)}{\partial x_i} \in H^1(\Omega), \quad \text{and} \quad \left\| \frac{\partial}{\partial x_i}(\chi u) \right\|_{H^1(\Omega)} \leq C \|f\|_{L^2(\Omega)}.$$

- It remains to prove that  $\frac{\partial^2}{\partial x_d^2}(\chi u) \in L^2(\Omega)$ : we re-use the original equation:

$$\frac{\partial^2}{\partial x_d^2}(\chi u) = g - \sum_{i=1}^{d-1} \frac{\partial^2}{\partial x_i^2}(\chi u).$$



**Proof of Step (iii).**

- By compactness of  $\bar{\Omega}$ , there exist open subsets  $\mathcal{O}_0 \Subset \Omega$ , and  $\mathcal{O}_1, \dots, \mathcal{O}_N \subset \mathbb{R}^d$  as in the statement of Step (ii) such that:


$$\bar{\Omega} \subset \bigcup_{i=0}^N \mathcal{O}_i.$$

- Let  $\{\theta_i\}_{i=0, \dots, N}$  be a **partition of unity** associated to the covering  $\{\mathcal{O}_i\}_{i=0, \dots, N}$ , i.e.

$$\forall i, \theta_i \in C_c^\infty(\mathcal{O}_i), \theta_i \geq 0, \text{ and } \sum_{i=0}^N \theta_i = 1 \text{ on } \bar{\Omega}.$$

- Then:

$$u = \underbrace{\theta_0 u}_{\substack{\in H^2(\Omega), \text{ by Step (i) and} \\ \|\theta_0 u\|_{H^2(\Omega)} \leq C \|f\|_{L^2(\Omega)}}} + \sum_{i=1}^N \underbrace{\theta_i u}_{\substack{\in H^2(\Omega), \text{ by Step (ii) and} \\ \|\theta_i u\|_{H^2(\Omega)} \leq C \|f\|_{L^2(\Omega)}}$$



## The Sobolev imbedding theorem

The **Sobolev imbedding theorem** states conditions for Sobolev class functions to be **regular** in the “classical” sense, i.e. for their belonging to a **Hölder space**  $C^{k,\sigma}(\Omega)$ :

$$u \in C^{k,\sigma}(\Omega) \Leftrightarrow \|u\|_{C^{k,\sigma}(\Omega)} := \|u\|_{C^k(\Omega)} + \sup_{|\alpha|=k} \sup_{\substack{x,y \in \Omega \\ x \neq y}} \frac{|\partial^\alpha u(x) - \partial^\alpha u(y)|}{|x - y|^\sigma} < \infty.$$

### Theorem 14 (Sobolev imbedding theorem).

Let  $\Omega \subset \mathbb{R}^d$  be a bounded Lipschitz domain. Let  $0 \leq k$ ,  $1 \leq m$  be two integers,  $1 \leq p < \infty$  be an exponent, such that there exists  $\sigma \in (0, 1)$  satisfying:

$$k + \sigma \leq m - \frac{d}{p}.$$





Then, the space  $W^{m,p}(\Omega)$  is continuously embedded in  $C^{k,\sigma}(\Omega)$ , and there exists a constant  $C > 0$  such that:

$$\forall u \in W^{m,p}(\Omega), \|u\|_{C^{k,\sigma}(\Omega)} \leq C \|u\|_{W^{m,p}(\Omega)}.$$






Roughly speaking, functions in  $W^{m,p}(\Omega)$  have “a little less” than  $m$  classical derivatives, and “tend to have  $m$  classical derivatives” as  $p \rightarrow \infty$ .

# Bibliography







## General mathematical references I

-  [All] G. Allaire, *Analyse Numérique et Optimisation*, Éditions de l'École Polytechnique, (2012).
-  [ErnGue] A. Ern and J.-L. Guermond, *Theory and Practice of Finite Elements*, Springer, (2004).
-  [EGar] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*, CRC Press, (1992).
-  [La] S. Lang, *Fundamentals of differential geometry*, Springer, (1991).

## Mathematical references around shape optimization I

-  [All] G. Allaire, *Conception optimale de structures*, Mathématiques & Applications, **58**, Springer Verlag, Heidelberg (2006).
-  [All2] G. Allaire, *Shape optimization by the homogenization method*, Springer Verlag, (2012).
-  [AlJouToa] G. Allaire and F. Jouve and A.M. Toader, *Structural optimization using shape sensitivity analysis and a level-set method*, J. Comput. Phys., 194 (2004) pp. 363–393.
-  [DelZol] M. Delfour and J.-P. Zolesio, *Shapes and geometries: metrics, analysis, differential calculus, and optimization*, Society for Industrial and Applied Mathematics, (2011).
-  [Ha] J. Hadamard, *Sur le problème d'analyse relatif à l'équilibre des plaques élastiques encastrées*, Mémoires présentés par différents savants à l'Académie des Sciences, 33, no 4, (1908).

## Mathematical references around shape optimization II






-  [HenPi] A. Henrot and M. Pierre, *Variation et optimisation de formes, une analyse géométrique*, Mathématiques et Applications 48, Springer, Heidelberg (2005).
-  [MoPir] B. Mohammadi et O. Pironneau, *Applied shape optimization for fluids*, 2nd edition, Oxford University Press, (2010).
-  [Mu] F. Murat, *Contre-exemples pour divers problèmes où le contrôle intervient dans les coefficients*, Annali di Matematica Pura ed Applicata, 112, 1, (1977), pp. 49–68.
-  [MuSi] F. Murat et J. Simon, *Sur le contrôle par un domaine géométrique*, Technical Report RR-76015, Laboratoire d'Analyse Numérique (1976).
-  [Pironneau] O. Pironneau, *Optimal Shape Design for Elliptic Systems*, Springer, (1984).
-  [Sethian] J.A. Sethian, *Level Set Methods and Fast Marching Methods : Evolving Interfaces in Computational Geometry, Fluid Mechanics, Computer Vision, and Materials Science*, Cambridge University Press, (1999).

## Mathematical references around shape optimization III



[SoZo] J. Sokolowski and J.-P. Zolesio, *Introduction to shape optimization*, Springer, Berlin, Heidelberg, (1992).

## Online resources I

-  [Allaire2] Grégoire Allaire's web page, <http://www.cmap.polytechnique.fr/allaire/>.
-  [Allaire3] G. Allaire, *Conception optimale de structures*, slides of the course (in English), available on the webpage of the author.
-  [AIPan] G. Allaire and O. Pantz, *Structural Optimization with FreeFem++*, *Struct. Multidiscip. Optim.*, 32, (2006), pp. 173–181.
-  [FreyPri] P. Frey and Y. Privat, *Aspects théoriques et numériques pour les fluides incompressibles - Partie II*, slides of the course (in French), available on the webpage <http://irma.math.unistra.fr/privat/cours/fluidesM2.php>.
-  [FreeFem++] *Web page of the FreeFem project*, <https://freefem.org/>.